

Alcohol and unprotected sex

Mixing sex with too much alcohol could leave you with more than you bargained for. Because alcohol can affect your judgment, it's easy to forget about using a condom or taking the pill which could leave you vulnerable to sexually transmitted infections, an unwanted pregnancy or a night you regret with someone that you're not even attracted to.

Facts and Figures

Sexually Transmitted Infections

The rapid spread of Sexually Transmitted Infections (STIs), is a significant health issue in the UK and has been shown to be related to alcohol consumption. The Government says that over 480,000 new cases were diagnosed in 2009. The infections are growing fastest among young people.⁽¹⁾

The two most common Sexually Transmitted Infections among young people are⁽²⁾:

Chlamydia: Because this STI sometimes shows no symptoms, it can be difficult to diagnose. It can cause an unpleasant discharge from the penis or vagina about two weeks after the infection occurs, and in the long term it can lead to chronic pelvic infection and infertility in women.

Genital warts: These hard, gritty bumps can appear around the genitals or anus. They don't always spread through penetrative sex - sometimes close genital contact is enough.⁽³⁾

More serious, but less common STIs, include syphilis, HIV and hepatitis B.

The best way to protect yourself against sexually transmitted infections is to wear a condom. But drinking can make you forget, or not bother in the first place. This is because alcohol can lower inhibitions and make you more likely to take risks⁽⁴⁾. In one survey, a third of 16-22 year olds said they have had unprotected sex when drunk.⁽⁵⁾ By ignoring advice on safe sex, or getting drunk, young people can run a serious risk of infection and unwanted pregnancy.

If they're not treated early, STIs can cause serious problems. But the good news is that many STIs are simple to treat if spotted early. So, if you think you might have one, make an appointment with your nearest Sexual Health/Genito-Urinary Medicine (GUM) clinic to get it sorted out. You can find your local clinic online at www.ruthinking.co.uk.

Alcohol and unwanted pregnancy

The other consequence of not using contraception is that you or your partner could get pregnant. The UK currently has one of the highest teenage pregnancy rates in Western Europe⁽⁶⁾. Not using a condom because you've been drinking could mean facing parenthood long before you feel ready.

If you think that you may be pregnant, help is available. You can get emergency contraception (sometimes called 'the morning after pill') free from most local pharmacies, as well as at a GUM or family planning clinic. It's effective up to 72 hours after you've had unprotected sex but the earlier you take it the better.

Alcohol and losing your virginity

Choosing to have sex for the first time can make you feel different emotions. You might feel excited, stressed, scared and nervous all at once.

Drinking to deal with these conflicting feelings may be an important factor when people lose their virginity. Since alcohol, like other drugs, lowers your inhibitions, drinking can seem like a good idea if you're nervous about being naked in front of another person, or worried about 'getting it right' in bed. But studies show that nearly one third of teens said they were unhappy with the way they lost their virginity, because they were too drunk⁽⁷⁾. It also doesn't help that too much alcohol can affect men's sexual performance.

Planning ahead is a good idea. If you think that you're likely to have sex, stay away from too much alcohol.




The 'beer goggles' effect

Too much alcohol can affect your sexual judgment⁽⁸⁾. Suddenly that bloke you don't really like becomes the best looking person at the party. This side effect of drinking can be very embarrassing. In a recent survey, nearly half (43%) of 10-17 year olds said they had kissed or slept with someone when drunk they wouldn't normally fancy.⁽⁹⁾

Going home with the wrong person can also be dangerous. In the same survey almost a quarter (22%) of people interviewed said they had gone home with a stranger, while 15% had invited a stranger home to their place.⁽¹⁰⁾ Being drunk makes you vulnerable, and getting intimate with someone you hardly know may lead to you going further than you want to go, or even end in rape and sexual assault.

Drinking and confidence

The results of a Drinkaware survey found that more than a third (39%) of 16 and 17 year olds drank alcohol to increase their confidence with the opposite sex⁽¹¹⁾. But, while a little 'Dutch courage' may appear to boost your confidence, the slurring and sickness that may come after too much alcohol can look less than attractive to the object of your affections.



Staying safe

Carry a condom: Whether you're planning to drink or have sex, a condom is the best way to protect yourself against STIs. It's a good idea to carry one just in case.

Stay in control: When it comes to sex, you are the person in control of what does or doesn't happen. You should never feel under pressure to go further than you're happy with.

Keep an eye out for your friends: If you see a friend leave with someone they don't know, call them over and make sure they're OK. They'll thank you for it later.

Stay the distance: Stick within the daily unit guidelines and try matching every alcoholic drink you have with something soft like water or fruit juice.

Eat before: Food can help slow down the absorption of alcohol into your bloodstream. Have a meal, like pasta or pizza, before you go out and eat a few snacks along the way to help you stay in control.

Where to go for further information and advice

Brook (www.brook.org.uk) is a national charity offering free and confidential sexual health advice and services to people under 25.

The Family Planning Association (www.fpa.org.uk) can help you make informed choices about sex and contraception.

For up-to-date interactive information and advice about sex and relationships visit R U Thinking? (www.ruthinking.co.uk).

Head to The Site (www.thesite.org) for regularly updated guides taking a comprehensive look at sexual health, alcohol, drugs and much more.

Drinkaware (www.drinkaware.co.uk) features the help and advice you need to make informed choices about sex and alcohol.

References

- 1 Cook, P.A. Et al., North West Public Health Observatory, 'Contributions of alcohol use to teenage pregnancy and sexually transmitted infection rates,' <http://www.cph.org.uk/showPublication.aspx?pubid=709>
- 2 Health Protection Agency 2010, 'Infection reports', Volume 4 Number 34, <http://www.hpa.org.uk/hpr/archives/2010/hpr3410.pdf>
- 3 Ibid. Health Protection Agency 2010
- 4 Lane, S 2004, 'Alcohol Effects on Human Risk Taking', Psychopharmacology, vol 172, no 1, pp 68-72.
- 5 Youthnet 2009, 'Sex Factor: Young People and Sexual Health'
- 6 Office for National Statistics 2010, 'Conceptions to women aged under 18 - annual numbers and rates', <http://www.statistics.gov.uk/statbase/Expodata/Spreadsheets/D8903.xls>
- 7 Independent Advisory Group on Sexual Health and HIV 2007, 'Sex, drugs and young people: A review of the impact drugs and alcohol has on young people's sexual behaviour'.
- 8 10 Abbey A et al 2000, 'Alcohol's Effects on Sexual Perception'. Journal of Studies on Alcohol vol 61, no 5, pp 688-697.

9 Youthnet 2009, 'Sex Factor: Young People and Sexual Health'.

10 Ibid. Youthnet 2009

11 YouGov plc online poll for Drinkaware, 3-6 July 2009. Total sample was 1,071 16-17 year olds living in Great Britain.

Contents approved by Drinkaware Chief Medical Adviser, Prof. Paul Wallace BSc (Hons), MSc, MBBS, FRCGP, FFPHM

Drinkaware

Samuel House
6 St Albans St
London SW1Y 4SQ
Registered in England & Wales No. 4547974.
A company limited by guarantee.
Registered Charity No. 1094586